


- 3 00000000000000


- 3 00000000000000




























01 | | | | |








分析某地人工林调查数据

(几种草本植物种群密度平均值,单位:株/m²)

0	0.2	0.4	0.6	0.8	1.0
15.3	13.5	10.8	7.4	4.3	2.4
10.4	9.5	6.1	5.6	3.0	1.2
3.7	4.3	8.5	4.4	2.2	1.0












0	0.2	0.4	0.6	0.8	1.0
15.3	13.5	10.8	7.4	4.3	2.4
10.4	9.5	6.1	5.6	3.0	1.2
3.7	4.3	8.5	4.4	2.2	1.0









1. []


01 | | | | | |














2. [













3. []



























02 生物因素







J形: 理想条件


S形:


随着种群的增长,<u>种内竞争</u>会加剧,从而使种群的增长<u>受到限制</u>, 这说明种群数量的变化受到种群内部生物因素的影响。P14





























5.双小核草履虫种群的K值为什么比单独培养时低?












条件	对于猞猁	猞猁种群出生率	猞猁种群死亡率













- ➤ TRH [] TSH [] [] []
- \succ 0000000000
- ▶ □□□□□□□□□□ TRH □ TSH □□□□□□"□"□






















		3 □/□
		4~6
	20%~30% 🛛	





























03 种群研究的应用











1. 00000000











2. ______








- 1
- K/2 00000000000000000
- **3"** [[[]"[]




3. 0000000


































()



- (1) DDDDDDDDDD (
 - **A.** 000000
 - **C.** ПППППППППП

- **B.** [[[[[]]]]]



- **D.**